



Sod-Seeding Small Grains Into Bermudagrass Pasture

Jeff Edwards

Small Grains Extension Specialist

Jason Warren

Soil and Water Conservation Extension Specialist

Daren Redfearn

Pasture and Forage Extension Specialist

When properly managed warm-season grasses, such as bermudagrass, are excellent forages and pasture crops for Oklahoma. Warm-season grasses, however, are dormant during the winter months, leaving producers with a need for supplemental pasture during this time. For this reason, the practice of sowing a small grain crop into established warm-season grasses (sod-seeding) is sometimes adopted in Oklahoma. Although the inputs required for this system are greater than warm-season pasture production alone, sod-seeded small grains can be a profitable system when managed properly and grain or beef prices are adequate.

Some reasons sod-seeding small grains into warm-season pasture makes sense for some Oklahoma producers include:

- *Reduced cash outlay for winter supplementation of livestock.* Protein supplements may not be needed when adequate small grain forage is available.
- *More efficient use of available land.* Warm-season grass pastures are out of production for about five months each year (November through April). A small grain crop can be grown during this period with minimal overlap.
- *Additional profits from grain production are possible.* Sod-seeded small grains are most frequently used for pasture only, but the potential also exists for grain production. However, more inputs are required and timing of inputs are more critical than in a forage-only or grain-only system.
- *More efficient use of seasonal precipitation in Eastern Oklahoma.* A significant portion of the precipitation received in eastern Oklahoma falls during the months of warm-season grass dormancy. A sod-seeded small grain crop, however, is actively growing during much of this time and can convert winter precipitation to forage more quickly than warm-season grasses alone.
- *Less erosion than conventionally-seeded small grains pasture.* Severe soil erosion problems can occur when soils are tilled each year for continuous small grains pasture. Implementing a sod-seeded small grain system on highly-erodible soils maintains cover, increases soil stability, and greatly reduces erosion potential.

Oklahoma Cooperative Extension Fact Sheets
are also available on our website at:
<http://osufacts.okstate.edu>



Sod-seeded small grains can provide forage during winter and early spring when warm-season grasses are dormant. In high-rainfall areas there is also the potential to harvest a grain crop, but both grain yield and forage yield is much less than in a monocrop system.

Adaptation

This system is best adapted to areas east of I-35 in Oklahoma. This is especially true when a forage and a grain crop are desired. The moisture requirement of small grains peaks during late April and May (heading to soft dough). If rainfall is not sufficient to replenish soil moisture following wheat harvest, yield of the subsequent hay or forage crop may be reduced. Since the probability of receiving adequate rainfall is greater in Central and Eastern Oklahoma, these regions are normally better suited to a sod-seeding system. Of course, the potential for excellent production with this system exists anywhere under irrigation.

Management

Site preparation

In most circumstances, a pre-plant application of a non-selective, contact herbicide such as paraquat (sold under various trade names including Gramoxone Extra) should be applied to temporarily terminate pasture growth and to eliminate weeds. A carrier of at least 15 gallons per acre of water is generally required to ensure adequate cover-

age. Paraquat is frequently the herbicide of choice, as it does not damage perennial grasses long-term. Systemic herbicides, such as glyphosate, have the potential to translocate to rooting portions of plants and can cause lasting damage. The other benefit of paraquat is that it does not require a significant amount of leaf area or new growth to ensure efficacy, so the warm-season grasses can be intensively grazed **prior** to herbicide application. As with all pesticides, there is the potential for drift and exposure, so be sure to read and follow all label recommendations as these can change from time to time.

Planting

Several planting methods can be used effectively for sod-seeding small grains pasture into established pastures. In all cases, the pasture top growth should be removed by grazing or mowing. This not only facilitates easier sowing, but also permits better sunlight availability to emerging small grain seedlings. A no-till drill in good working order is the best method to ensure good seed-to-soil contact and uniform emergence. Ideally, a drill used for sowing small grains should 1) have a row spacing of 7.5 inches or narrower, 2) provide enough down force to ensure accurate seed placement in hard soils, and 3) have closing wheels linked to the disk opener(s).

A no-till drill is not essential to establishing small grains in sod. Some producers have obtained acceptable plant stands using a conventional-drill with double disk openers. Acceptable plant stands can also be obtained by disking and then broadcasting seed, but it is important to note that this method will result in the most inconsistent seeding depth. Ultimately, there are several methods that can be used to obtain a stand of small grains sufficient for grazing or hay production. If grain production is the end goal, however, a more uniform, consistent stand is required and the use of a grain drill is advised.

Small grains choice

While wheat is not the only option for small grains pasture, it is the most popular in Oklahoma. Rye is usually the best forage producer in late fall and winter because it grows at cooler temperatures than other small grains. It performs very well when sod-seeded on sandy soils. Oat is an excellent spring forage producer, but may winterkill if sown in the fall in the northern half of the state. Oat and barley may also be sown in the spring (February) for grain or forage production.

If wheat is to be sown, there are several factors to consider. Variety selection is very important if a grain crop is to be harvested, but less important if forage-only production is the goal. Varieties differ greatly in their ability to recover from grazing and produce a grain crop. Consult a recent copy of the OSU Small Grains Variety Performance Tests (available at www.wheat.okstate.edu) to determine which wheat varieties recover well from grazing. If grain is to be harvested, it is also extremely important to plant weed-free seed. Weeds are very difficult to control in a sod-seeded system, and introducing additional weeds at planting only makes the job harder.

If forage or hay is the primary objective, then factors such as planting date and seeding rate are far more influential than variety in determining forage production. Increasing the seeding rate from 60 lb/acre to 120 lb/acre, for example, results in adequate forage production by most



The increased use of no-till planting methods in grain cropping systems, makes high-quality no-till grain drills, such as the one shown above, readily available in the used equipment market.

commercially available wheat cultivars. It is also important to note that most commercially available wheat cultivars are now covered by the Plant Variety Protection Act (PVP) and can only be sold as a certified class of seed. While farmers can still save and plant seed of PVP-protected wheat varieties on their own farm, purchase and planting of 'bin-run' seed of a PVP-protected variety is illegal.

Planting date

The optimum planting date for sod-seeded small grains depends on which small grain is chosen and the method chosen to suppress pasture-grass competition and control weeds prior to planting. If paraquat is used, planting could begin as early September. The recovery time for bermudagrass after paraquat application is about two to four weeks, depending on how early it is applied and temperatures. This is usually long enough for small grains to become established. If chemical control is not used, the warm-season grasses should be grazed close to the ground and disked. If this option is used, planting should be delayed until October. Remember October planting may reduce fall forage production.

Fertilization

A soil test should be taken prior to seeding small grains, and any pH problems should be corrected prior to seeding. In most cases, nitrogen is required to meet crop needs. If forage is the primary objective, nitrogen should be applied pre-plant or in early winter. If forage plus grain is the end goal, it is advisable to split the nitrogen application between an early winter application and a second application after cattle have been removed. The amount of nitrogen applied depends on the yield goal of forage or grain. When determining yield goals, assume it takes 30 to 40 lbs N/ acre to produce 1,000 lbs of forage and 2 lbs N/ acre to produce one bushel of grain. The yield goal should be based on past experience and in-season assessment of yield potential.

Phosphorous (P) is especially important in the early stages of small grain growth. The most efficient way to provide it to

seedlings is an in-furrow band. Banding P fertilizer into the row with seed presents little danger to seedlings; however, the amount of nitrogen plus potassium applied with the seed should not exceed 30 lbs/acre total.

Pest control

If forage is the primary objective, in-season weed control may not be necessary. If grain is to be harvested, it is necessary to control winter annual weeds with a herbicide. Many herbicides labeled for wheat have grazing restrictions and some do not control all winter annual weeds present in Oklahoma wheat fields. So, it is extremely important to know what weeds are present and to read and follow label directions carefully. In addition to herbicides, insecticides may be required to control insect pests such as fall armyworm, grasshoppers, or cutworms.

Expectations

A properly fertilized sod-seeded grain crop can provide 0.75 to 1.5 tons/acre of fall forage and two to three tons/acre of full-season forage. To achieve this type of production, however, 60 to 150 lb/acre of nitrogen is required and the bulk of this should be provided in the form of nitrogen fertilizer. A grain yield of 20 to 30 bu/acre can be achieved in a dual-use system if grazing is terminated at the first hollow stem stage of growth, weeds are managed, and adequate fertility is supplied. A yield goal of 1 ton per acre of forage followed by 20 bu/acre of grain, for example, would require 100 lb/acre of nitrogen. So, while sod-seeded small grains pasture can provide supplemental grazing and grain income, it should not be viewed as a cheap source of additional grazing.

The Oklahoma Cooperative Extension Service

Bringing the University to You!

The Cooperative Extension Service is the largest, most successful informal educational organization in the world. It is a nationwide system funded and guided by a partnership of federal, state, and local governments that delivers information to help people help themselves through the land-grant university system.

Extension carries out programs in the broad categories of agriculture, natural resources and environment; family and consumer sciences; 4-H and other youth; and community resource development. Extension staff members live and work among the people they serve to help stimulate and educate Americans to plan ahead and cope with their problems.

Some characteristics of the Cooperative Extension system are:

- The federal, state, and local governments cooperatively share in its financial support and program direction.
- It is administered by the land-grant university as designated by the state legislature through an Extension director.
- Extension programs are nonpolitical, objective, and research-based information.
- It provides practical, problem-oriented education for people of all ages. It is designated to take the knowledge of the university to those persons who do not or cannot participate in the formal classroom instruction of the university.
- It utilizes research from university, government, and other sources to help people make their own decisions.
- More than a million volunteers help multiply the impact of the Extension professional staff.
- It dispenses no funds to the public.
- It is not a regulatory agency, but it does inform people of regulations and of their options in meeting them.
- Local programs are developed and carried out in full recognition of national problems and goals.
- The Extension staff educates people through personal contacts, meetings, demonstrations, and the mass media.
- Extension has the built-in flexibility to adjust its programs and subject matter to meet new needs. Activities shift from year to year as citizen groups and Extension workers close to the problems advise changes.

Oklahoma State University, in compliance with Title VI and VII of the Civil Rights Act of 1964, Executive Order 11246 as amended, Title IX of the Education Amendments of 1972, Americans with Disabilities Act of 1990, and other federal laws and regulations, does not discriminate on the basis of race, color, national origin, gender, age, religion, disability, or status as a veteran in any of its policies, practices, or procedures. This includes but is not limited to admissions, employment, financial aid, and educational services.

Issued in furtherance of Cooperative Extension work, acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture, Robert E. Whitson, Director of Cooperative Extension Service, Oklahoma State University, Stillwater, Oklahoma. This publication is printed and issued by Oklahoma State University as authorized by the Vice President, Dean, and Director of the Division of Agricultural Sciences and Natural Resources and has been prepared and distributed at a cost of 20 cents per copy. 0409 GH